



Review Article

Different applications of laser in the field of orthodontics: A review

Rashi Chaturvedi^{1,*}, Syed Tousif Ulla¹, Farhan Hussain¹, Zahid Manzoor¹,
Rahul Vaid², Nasir Ul Sadiq¹

¹Dept. of Orthodontics, Kalka Dental College, Meerut, Uttar Pradesh, India

²Periodontist and Implantologist, Meerut, Uttar Pradesh, India



ARTICLE INFO

Article history:

Received 01-04-2021

Accepted 22-04-2021

Available online 12-07-2021

Keywords:

Lasers

tooth movement

pain

reducing pain threshold

lasers in orthodontics

ABSTRACT

In today's time laser has been used widely in various fields of dentistry, out of all one is in the field of orthodontics and proved to be as useful equipment. Specific lasers such as soft tissue lasers can be used in the treatment of exposure of the tooth surgically with its prime advantage of very less bleeding as well as little bit of swelling it is also helpful in reducing the potential of post reducing pain. Along with all this the other applications of lasers in the field of orthodontics are, they are helpful in etching the superior most surface of the tooth i.e. enamel as well as helpful in bonding of the bracket to the enamel and vice versa i.e. debonding of the bracket from the superior most surface of the tooth i.e. enamel. Lasers are also helpful in tooth movement along with reducing the pain threshold.

© This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Introduction

Term laser denotes “light amplification by stimulated emission of radiation”. A device which is capable of transforming electrical form of energy in to the optical form of energy and results in light formation is known as a laser. Lasers produced monochromatic light of beam and which is coherent and is absolutely opposite to the lights which are ordinary.¹ The basic concept of lasers were derived from the quantum theory of radiation which was given by sir Einstein.² The maser technique which was studied by towns ultimately resulted in infrared along with optical region, which ultimately resulted in the invention of lasers.³ and the first laser was developed by maiman in the year 1960.⁴ lasers were introduced in the field of medical by goldman.⁵

The first effect of lasers over the dental caries were demonstrated by goldman in the year 1964, and revealed that lasers are helpful equipment in the reduction of dental caries.⁶ When the enamel is exposed to ruby laser energy tooth respond in different manners like there are

crater formation along with dentine fusion along with disappearance of dental caries, some other authors respond with the same results.^{7,8} laser technique were extended to both the hard and soft tissues in the oral cavity with the invention of neodymium doped yttrium aluminum garnet (Nd:YAG) laser and carbon dioxide (CO₂) laser in the year of 1964.⁹ in the year of 1980 Nd:YAG laser was used first timely in the oral cavity, for the prevention of dental caries by the researcher. Between the year 1970 and 1980 various researches over the different applications CO₂ lasers on the hard and soft tissue were tested.^{9,10} After 1990 various newer types of lasers were invented and used over hard and soft tissue in the oral cavity like Holmium: YAG, Erbium:YAG, Argon, Erbium yttrium scandium gallium garnet.

The basic theory quantum of physics were firstly described by neils bohr.¹¹ the smallest unit of energy i.e. photon is released after another photon is absorbed by the atom. This is known as spontaneous emission phenomenon. This theory were further developed by sir Einstein. He added to the basic theory that, a single excited atom is being able to absorb energy quantum and then released

* Corresponding author.

E-mail address: dr.rashichaturvedi@gmail.com (R. Chaturvedi).

two photons instead of one which result in formation of a single coherent wave in two photons travel. By these identical photons more atoms will get excited and results in emission of further identical photons, which ultimately results in light energy that is amplified totally and finally results in production of laser beam.¹²

1.1. Different properties of laser

1. **Coherency** : Stated that all the different types of light waves are identical in laser light.
2. **Collimation** : The beam which is present in a laser light should have a unique or straight direction as well as size and shape when compared to the conventional light, which gets diverged in all the direction.
3. **Monochromaticity** : The light which is produced by a laser is of narrow wave length when compared to the conventional light. That's why laser light is having single color.

The laser is basically made up of three components i.e. resonant optical, which is made up of mirror, semi-conductor which act as an active medium and the most importantly the energy source.

There are basically two different types of laser beam delivery system,

1. Interior mirror with hollow tube like wave guide: In this the beam of laser is being reflected by the mirrors along the wave guide and exit through the hand piece. In this there is no direct contact of the laser with the tissue, basically carbon dioxide type of laser used this technique.
2. Optic cable along with glass fiber: This cable is available in different sizes ranging from 200 microns to 1000 microns. It is enclosed in a sheath and is very fragile and it can't be bent to any sharp angle. This type of laser can be used in both the types i.e. with the direct tissue contact and without the direct tissue contact, this type of laser is used most commonly in direct tissue contact type fashion.

Light energy is being emitted by the laser in three different modes, i) when the laser beam is emitted in continuous power level it is known as ii) in second mode there is alteration in the power of the energy like on and of types and is known as gate pulsed mode. Iii) And the last type is known as free running pulsed mode in which major amount of energy is released for short duration of time.

1.2. Effect of laser

Whenever the laser came in contact with the oral tissue, for different types of interactions occur between the laser and the oral tissue. When there occur no effect over the target site and the laser beam gets reflected away from the tissue

site, known as reflection. Second type is, when the laser beam or energy is transmitted through the tissue and show no effect in the target site or the target tissue is known as transmission. The third is when heat being produced by the laser site and is transferred to the adjacent site is known as scattering. The most important and prime effect of laser is when it hits the target site or the target tissue in the oral cavity and produced its effect over it is known as the absorption of the laser energy.

2. Different applications of lasers in the field of orthodontics

Different types of abnormalities related to soft tissue might occur during the time or before or after the orthodontic treatment. Basically there are three different types conditions that may occur with respect to orthodontic treatment i.e. abnormal growth of the gingival soft tissue,¹³ abnormal attachment of frenum, or impacted tooth.^{14–18} Minor surgical procedures such as gingivectomy, frenectomy, can be performed with the help of lasers. Lasers can also be used in surgical exposure of the tooth. Diode, CO2 and Nd:YAG, are the most commonly used one lasers but recently Er:YAG and Er,Cr:YAG achieved more attention by their use in the field of soft tissue surgery.^{19–21} Clinicians found difficulty in placing the brackets in enlarged or irregularly contoured gingiva which may lead to change or alter the height of the crown which may alter the tooth proportion.²² So for the gingivectomy or gingivoplasty procedure carbon dioxide lasers are used most commonly because of its wavelength is well absorbed by the soft tissue and is composed of water. After the year of 1980 these carbon dioxide based lasers are being used by periodontists to remove the irregularly shaped or enlarged gingiva for aesthetic reason as well as for the functional reason.^{23,24}

Another laser known as Nd:YAG and diode lasers were also used in as a treatment modality for the procedure of frenectomy and clinicians found satisfactory results with the same lasers when compared to carbon dioxide lasers.^{25,26} These lasers work on the principle of continuous or interrupted continuous wave mode. The dental hard tissues are not being able to well absorb the wave length of the laser. The only possible drawback of this type of laser is production of thermal heat and this could be the only possible reason that this type of lasers cannot be used in direct contact with the hard tissue i.e. alveolar bon.²⁷

Literature revealed that Er:YAG laser induced mild analgesic effect.²⁸ This laser works on a principle of pulse waved manner and simultaneously absorbed well by dental hard tissue as well as water and it results in minimal thermal effect over the hard tissues and is well tolerated. This laser can be used in conjunction with Nd:YAG or diode lasers for the process of frenectomy.

Laser technique can be used in the surgical exposure of impacted tooth at the bone level or at mucogingival level. When the exposure of the impacted tooth is by the help of lasers, it is clean, painless, and very quick. A totally dry field is achieved when the tooth is exposed by laser, and this could help in easy placement of bracket over the exposed tooth surface.²⁹ However literature revealed the occurrence the sub cutaneous emphysema after the tooth is surgically exposed with the help of lasers.³⁰ Erbium laser gained popularity in the field of surgical exposure of impacted tooth because it is well tolerated by hard and soft tissues. This laser is also helpful in etching the outer surface of the tooth i.e. the enamel and facilitate in bonding of the orthodontic bracket to the tooth surface.³¹ The bond strength of the tooth i.e. etched via laser is found to be same to other tooth surface that is etched with the help of acid.³² The phosphate to calcium ratio of the enamel can be altered after irradiation with the use of lasers which ultimately leads in formation of stable acid proof compound.³²

With the use of ceramic brackets in orthodontics there are higher chances of fracture of enamel because of high bond strength of the ceramic brackets. It can be overcome by the process of laser irradiation, it softens the resins when the brackets are heated and thus helps in reducing the force required for debonding. Basically two principles are used in laser debonding i.e. photoablation and thermal softening.^{33,34} Literature revealed that low level laser therapy is very much helpful in reducing in pain threshold in tooth movement cases in first 5 to 7 days of the tooth movement. Some investigated pain relief with CO2 laser after the first wire was placed, while others studied pain relief after separators placement with gallium-aluminum arsenium and CO2 lasers.³⁵

Low-level lasers can accelerate the opening of the mid-palatal suture and improve bone regeneration during and after rapid maxillary expansion according to several studies.

Safety from dental laser is very much required, laser can affect retinal damage can cause cataract, so one should wear proper eye protection from the wavelength of the laser. Lasers can cause blistering and burning sensation to the skin, the clinician should cover the full body while working with lasers. Lasers may also cause coughing, nasal congestion by emission of fumes containing noxious elements. So the clinician should wear masks and high volume evacuation during the use of lasers.

3. Conclusion

Lasers are becoming very useful in the treatment modality of orthodontics. Use of lasers provide good hemostasis, reduced post operating pain as well as swelling enhanced wound healing and total control over the incision. One should follow strict rules while using the lasers.

4. Source of Funding

No financial support was received for the work within this manuscript.

5. Conflicts of Interest

There are no conflicts of interest.

References

- Gould RG. The LASER, Light Amplification by Stimulated Emission of Radiation. The Ann Arbor Conference on Optical Pumping, the University of Michigan, June 15 through June 18; 1959.
- Einstein A. Zur Quantum Theorie der Strahlung. *Phys Z.* 1917;18:121-8.
- Schawlow AL, Townes CH. Infrared and Optical Masers. *Phys Rev.* 1940;112(6). doi:10.1103/PhysRev.112.1940.
- Maiman T. Stimulated Optical Radiation in Ruby. *Nature.* 1960;187(4736):493-4. doi:10.1038/187493a0.
- Goldman L, Blaney DJ, Kindel DJ, Richfield DF, Owens P, Homan EL, et al. Effect of the Laser Beam on the Skin. *J Invest Dermatol.* 1963;40:121-2.
- Goldman L, Hornby P, Meyer R, Goldman B. Impact of the Laser on Dental Caries. *Nature.* 1964;203(4943):417. doi:10.1038/203417a0.
- Goldman L, Gray JA, Goldman B, Meyer R. Effects of laser: impact on teeth. *J Am Dent Assoc.* 1965;70:601-6.
- Stern RH, Sognnaes RF. Laser effect on dental hard tissues. A preliminary report. *J South Calif Dent Assoc.* 1965;33:17-9.
- Pecaro BC, Garehime WJ. The CO2 laser in oral and maxillofacial surgery. *J Oral Maxillofac Surg.* 1983;41(11):725-8. doi:10.1016/0278-2391(83)90189-1.
- Frame JW. Carbon dioxide laser surgery for benign oral lesions. *Br Dent J.* 1985;158(4):125-8. doi:10.1038/sj.bdj.4805545.
- Bohr N. The theory of spectra and atomic constitution. In: 2nd Edn. Cambridge: Cambridge University Press; 1922.
- Coluzzi DJ. Fundamentals of dental lasers: science and instruments. *Dent Clin North Am.* 2004;48(4):751-70. doi:10.1016/j.cden.2004.05.003.
- Ruffoli R, Giambelluca MA, Scavuzzo MC, Bonfigli D, Cristofani R, Gabriele M, et al. Ankyloglossia: a morphofunctional investigation in children. *Oral Dis.* 2005;11(3):170-4. doi:10.1111/j.1601-0825.2005.01108.x.
- Baer PN, Cocco PJ. Case Report: Gingival Enlargement Coincident with Orthodontic Therapy. Report of Three Cases. *J Periodontol.* 1964;35(5):436-9. doi:10.1902/jop.1964.35.5.436.
- Zachrisson S, Zachrisson B. Gingival condition associated with orthodontic treatment. *Angle Orthod.* 1972;42(1):26-34.
- Costuleanu M, Plamadalea P, Costuleanu N, Foia L, Indrei A. The causes of gingival overgrowth. *Rev Med Chir Soc Med Nat Iasi.* 1999;103:63-7.
- Kouraki E, Bissada NF, Palomo JM, Ficara AJ. Gingival enlargement and resolution during and after orthodontic treatment. *N Y State Dent. N Y State Dent J.* 2005;71:34-7.
- Huang WJ, Creath CJ. The midline diastema: A review of its etiology and treatment. *Pediat Dent.* 1995;17:171-9.
- Cobb CM, Low SB, Coluzzi DJ. Lasers and the Treatment of Chronic Periodontitis. *Dent Clin North Am.* 2010;54(1):35-53. doi:10.1016/j.cden.2009.08.007.
- Barak S, Kaplan I. The CO2 laser in the excision of gingival hyperplasia caused by nifedipine. *J Clin Periodontol.* 1988;15(10):633-5. doi:10.1111/j.1600-051x.1988.tb02264.x.
- Gontijo I, Navarro RS, Haypek P. The applications of diode and Er:YAG lasers in labial frenectomy in infant patients. *J Dent Child.* 2005;72(1):10-5.
- Sarver DM, Yanosky M. Principles of cosmetic dentistry in orthodontics: Part 2. Soft tissue laser technology and cosmetic gingival contouring. *Am J Orthod Dentofac Orthop.* 2005;127(1):85-90. doi:10.1016/j.ajodo.2004.07.035.

23. Abt E. CO2 laser treatment for gingivectomies reduces hemorrhaging, post-op pain. *Clin Laser Mon.* 1992;10(1):8–8.
24. Gama S, de Araújo T, Pinheiro A. Benefits of the use of the CO2 laser in orthodontics. *Lasers Med Sci.* 2008;23(4):459–65. doi:10.1007/s10103-007-0506-2.
25. Abdel-Aziem F. Clinical evaluation of pulsed ND:YAG dental laser applied on oral soft tissues. *Egypt Dent J.* 1994;40(4):863–70.
26. Kara C. Evaluation of patient perceptions of frenectomy: a comparison of Nd:YAG laser and conventional techniques. *Photomed Laser Surg.* 2008;26(2):147–52.
27. Sarver DM. Use of the 810 nm diode laser: soft tissue management and orthodontic applications of innovative technology. *Pract Proced Aesthet Dent.* 2006;18:7–13.
28. Aras MH, Göregen M, Güngörmüş M, Akgül HM. Comparison of Diode Laser and Er:YAG Lasers in the Treatment of Ankyloglossia. *Photomed Laser Surg.* 2010;28(2):173–7. doi:10.1089/pho.2009.2498.
29. Juan RB, Miguel H, Enrique E, Clervie P, Antonio E. Erbium Laser Treatment Of An Impacted First Mandibular Premolar: A Case Report. *J ClinPediatr Dent.* 2008;33(1):9–12. doi:10.17796/jcpd.33.1.6867w5q5r71p6167.
30. Dabiri D, Nedley M, Baugh R, Haerian A. An incident of subcutaneous emphysema following laser tooth exposure. *J Clin Orthod.* 2012;46(12):749–52.
31. Genovese MD, Olivi G. Use of laser technology in orthodontics: hard and soft tissue laser treatments. *Eur J Paediatr Dent.* 2010;11(1):44–8.
32. Klein ALL, Rodrigues LKA, Eduardo CP, dos Santos M, Cury JA. Caries inhibition around composite restorations by pulsed carbon dioxide laser application. *Eur J Oral Sci.* 2005;113(3):239–44. doi:10.1111/j.1600-0722.2005.00212.x.
33. Ghafari J, Skanchy TL, Mante F. Shear bond strengths of two ceramic brackets. *J Clin Orthod.* 1992;26(8):491–3.
34. Bishara SE, Fonseca JM, Boyer DB. The use of debonding pliers in the removal of ceramic brackets: Force levels and enamel cracks. *Am J Orthod Dentofacial Orthop.* 1995;108(3):242–8. doi:10.1016/s0889-5406(95)70016-1.
35. Eslamian L, Borzabadi-Farahani A, Hassanzadeh-Azhiri A, Badiee MR, Fekrazad R. The effect of 810-nm low-level laser therapy on pain caused by orthodontic elastomeric separators. *Lasers Med Sci.* 2014;29(2):559–64. doi:10.1007/s10103-012-1258-1.

Author biography

Rashi Chaturvedi, 3rd Year MDS Student

Syed Tousif Ulla, 2nd Year MDS Student

Farhan Hussain, 1st Year MDS Student

Zahid Manzoor, 1st Year MDS Student

Rahul Vaid, Clinician

Nasir Ul Sadiq, 2nd Year MDS Student

Cite this article: Chaturvedi R, Ulla ST, Hussain F, Manzoor Z, Vaid R, Sadiq NU. Different applications of laser in the field of orthodontics: A review. *IP Indian J Orthod Dentofacial Res* 2021;7(2):90-93.