

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Indian Journal of Orthodontics and Dentofacial Research

Journal homepage: <https://www.ijodr.com/>

Original Research Article

The effects of erupting mandibular 3rd molar on the dental arch: A FEM studyAvinash Kumar^{1,*}, Meghna Bhandhari¹, MD Baba Fareeduddin²¹Dept. of Orthodontics & Dentofacial Research, Rajiv Gandhi University of health sciences, Kalaburgi, Karnataka, India²Dept. of Orthodontics, ESIC Dental College and Hospital, Gulbarga, Karnataka, India

ARTICLE INFO

Article history:

Received 11-06-2022

Accepted 13-07-2022

Available online 27-10-2022

Keywords:

Mesio-angular impaction

Horizontal impaction

Finite element study

Stress distribution

Tooth displacement.

ABSTRACT

Introduction: Role of Erupting Mandibular 3rd molars as a cause of incisor crowding in the lower arch continues to be controversial. The relation between 3rd molar and dental crowding has not been established.

Aims and Objectives: To determine the area of stress distribution and tooth displacement during eruption of mandibular 3rd molar in mesioangular and horizontal impaction.

Materials and Methods: Three dimensional finite element models were generated based on computed tomography scan data. From computed tomographic scans of a human mandible two additional finite element models were generated: a mandibular model with mesioangular impaction and second model with horizontal impaction. To investigate the stress distribution to the mandible, and tooth displacement eruptive force of 10 grams was applied.

Results: High concentration of stresses was seen at the inferior border of the mandible and lowest concentration of stress was present at symphysis for both the models. The tooth displacement for both mesio-angular and horizontal impaction caused mesial tilting of all the teeth i.e. from 2nd molar to central incisor; with greatest tilting of 2nd molar. Along the vertical plane, there was extrusion seen with 2nd molar with mesio-angular impaction and extrusion in 2nd molar and 1st molar with horizontal impaction. And all the teeth exhibited buccal flaring in case of both impaction.

Conclusion: There is significant level of tooth displacement from 2nd molar towards the incisors. These changes occurring along the dental arch must be taken into consideration while planning for prophylactic extraction of third molars.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Lower arch crowding is an ongoing source of controversy and concern in orthodontics. The role of 3rd molars as a cause of incisor crowding, especially in the lower arch continues to be controversial. From the early beginning of orthodontics as a specialty, the presence of 3rd molars was believed by orthodontists to be responsible for late crowding of the lower anterior teeth. Because this was frequently observed coincident with the timing of 3rd molar eruption,

clinicians were tempted to conclude a “cause and effect” relationship between these two phenomena. At a simplistic level, it was reasoned that vector forces from erupting 3rd molars were pushed against second molars, causing mesial migration of the posterior teeth. The result was loss of available space and crowding. The relation between 3rd molar and dental crowding has not yet been established. Clinicians have always been divided between supporters and opponents of anterior dental crowding caused due to the force generated by the 3rd molar eruption. For the same reason, the surgical prophylactic approach for 3rd molar has

* Corresponding author.

E-mail address: dr.avinashsajjan@gmail.com (A. Kumar).

always been seen as the cure by the former and a 'placebo' by the latter.¹

Late crowding of the lower incisor teeth is frequently observed after the eruption of the 3rd molars, inciting the clinicians to deliberate a cause-and-effect relationship between the two events. The hypothesis is presumed that the mesial component of the forces created by the erupting 3rd molars, transmitted through the dental arch, can create a mesial migration of the teeth culminating in the area of the incisors. The result is the inaugural loss of available space and crowding. Lower arch crowding that develops or increases after establishment of the permanent dentition during the teenage period, can be best described as post-adolescent crowding. The role of erupting 3rd molars as a compel of such dental crowding has been the subject of controversy over the years.

Various theories have been proposed that account for late lower arch crowding which includes late mandibular growth, anterior component of force, lack of attrition in the modern population which may possibly cause mesial tilting of these teeth. This mesial axial inclination of permanent teeth gives rise to anterior component of force. Anterior component of force is an occlusal force dissipated axially and in the mesial direction, i.e. towards the frontal region of the mouth. This force is transmitted through the proximal tooth contacts of teeth, anterior to the mesially tilted 3rd molars. This force produced by 3rd molar has a possible role in mesial migration of teeth anterior to it. Subsequently, this might be the reason of malalignment of anterior teeth and crowding. In the year 1961, Bergstorm² was among one of the first authors to analyze the influence of 3rd molars in the developing dental arch crowding and reported that there was a relationship between wisdom teeth and incisor crowding. Bishara et al³ evaluated changes in the dental arches between 25 and 45 years of age and concluded that both sexes experienced a significant increase in dental crowding in both the dental arches, however the severity was more pronounced in the lower arch especially in the lower anterior segment. Later, Vego's⁴ study concluded that the eruption of lower 3rd molars exerted a force on neighbouring teeth as well. On the contrary to this, Broadbent⁵ supported the opposite theory, that the presence of 3rd molars had no influence on crowding. Sidlauskas⁶ also reported that there is no co-relation between 3rd molar and anterior crowding.

The finite element analysis (FEA) is an upcoming and significant research tool for biomechanical analyses in biological research. It is an ultimate method for modeling complex structures and analyzing their mechanical properties. FEA has now become widely accepted as a non-invasive and excellent tool for studying the biomechanics and the influence of mechanical forces on the biological systems. It enables the visualization of superimposed structures, and the stipulation of the material properties of anatomic craniofacial structures. It also allows in

establishing the location, magnitude, and direction of an applied force, as it may also assign stress points that can be theoretically measured. Further, as it does not affect the physical properties of the analyzed materials it is easily repeatable. Thus, finite element-three dimensional analysis gives more accurate, adaptable and clear picture of the areas of tooth displacement and stress effects of erupting 3rd molars on lower anterior crowding. Thus, the study aims in determining the areas of stress distribution and tooth displacement caused by erupting 3rd molar.

2. Materials and Methods

FEA has been developed into a branch of applied mathematics for numeric modelling of physical systems, which is used in many engineering disciplines. In its simplest mathematical terms, this numerical technique is used to find approximate solutions for partial differential and integral equations through the generation of meshes of a continuous domain for a set of discrete sub domains or elements. Numerical methods are then used to predict the behaviour of the object in question in various situations, for example, under conditions of loading.⁷

The external forces and the mechanical properties/geometry are used to calculate the nodal displacements; the differentiation of the displacement field yields the strain distribution; and the stress distribution is determined mathematically.

The mandibular model was developed from a patient without any gross defects or discontinuity in the anatomy. The model had a set of permanent dentition with normally erupted teeth, without any crowding or spacing and impacted 3rd molar on one side and a normally erupted 3rd molar on the other side. Computed tomographic (CT) scan images of the mandible were made in the axial direction at 0.5mm intervals in the horizontal plane. The study aimed to determine the stresses and displacements exerted by the mandibular 3rd molar.

In this study, CT scan of the mandible will be used to build a 3 – dimensional FEA model. There were 2 finite element models constructed, the first one with mesio-angular impaction and the other with a horizontal impaction. As mentioned in the literature, the eruptive force of tooth was found to be between 5-10 grams,⁸⁻¹² the models using 10 grams of force will be taken into account for determining the areas of maximum stress and tooth displacement. The software used for geometric modeling was Solid Works. Individual CT scan sections will be imported into the Solidworks and traced, taking care so as not to distort the anatomy of the region. Each traced sections will be then placed one above the other to give a mandibular model in Solidworks. These blocks will be imported into Altair HyperMesh. The software which will be made use in present study is Hyperworks 13.0, Altair, Troy, Michigan, United States.

2.1. Generation of CAD Model

The geometric model obtained is converted to finite element model using Altair HyperMesh software. FEM has been applied successfully to study stress and strain in the field of engineering and in living structure. The FEM is composed of an aggregate of small elements that are sufficient to describe the geometry of the subjects. This is called ‘creating the mesh or meshing’. The mesh intersections are called nodes. The contour data of the profiles created by surface scanning were transformed into the x, y and z co-ordinate points and read by Finite element program HyperWorks. Connecting these coordinate points gave line geometry also called as wire frame modeling. Connecting the lines of each section gave surface geometry.

2.2. Generation of finite element model

The finite element modelling is the representative of geometry in terms of finite number of elements and nodes. This process is called discretization. The main idea behind discretization is to improve the accuracy of the results. Subdivide i.e. discretize the complex geometry into suitable set of smaller “elements” of finite dimensions (2D or 3D). The points connecting two or more discrete elements are called as nodes or nodal points. The corner nodes are called primary external nodes. The additional nodes which occur on the sides of the elements are called secondary external nodes. The secondary nodes have fewer displacements than corner nodes. For this study Tetrahedron elements were used. Images of the Mesh are as shown (Figure 1).

Number, size and type of element are decided. Practical knowledge and judgement are needed to limit the number of elements to minimum amount conducive to acceptable results (Table 1). The material properties of the bone and teeth in the model were defined according to experimental data from previous studies (Table 2).

The boundary conditions are defined to simulate how the model is constrained and to prevent it from free body motion. Restraints will be established at all nodes lying at the lower border of the mandibular bone lying on the symmetric plane and appropriate boundary conditions are imposed.

2.3. Loading configuration

For determining the changes, 10 grams of force in both the horizontal and mesio-angular impaction of mandibular 3rd molar will be used.

3. Results

3.1. Stress distribution

When the eruptive forces were applied on the mandible, the stresses were observed at symphysis, body of the mandible, inferior border of the mandible, anterior border of ramus and the retro molar area in both models. Even though the

stresses were generally distributed throughout the mandible, a high concentration of stresses was present at the inferior border of the mandible and lowest concentration of stress was present at symphysis for both the models. The stresses were also concentrated in the body of the mandible, inferior border of the mandible, anterior border of ramus and the retro molar area (Table 3).




Fig. 1: a: FEM model 1- Mesioangular impaction; b: FEM model 2- Horizontal impaction.




Fig. 2: Stress distribution in the mandible – Buccal view; a: Mesio-angular impaction; b: Horizontal impaction.




Fig. 3: Stress distribution in the mandible – Lingual view. a) Mesio-angular impaction. b) Horizontal impaction.

Table 1: Meshing details

Model	Mesioangular.		Horizontal.	
	No. of nodes.	No. of elements.	No. of nodes.	No. of elements.
Teeth.	23287	107429	23287	107429
Periodontal Ligament.	69774	104827	69824	106443
Alveolar bone.	55622	286750	56551	292497

Table 2: Material property data representation.

Material	Young’s Modulus (Newton/Mm2)	Poisson’s Ratio
Teeth	20.3 × 103	0.30
Compact bone	13.7 × 103	0.30
Cancellous bone	79 × 102	0.30
Periodontal ligament	68.9	0.45

Table 3: Tabular presentation of stress distribution in mandible.

Areas of Mandible.	Mesioangular Impaction.	Horizontal Impaction.
Symphysis.	2.0E-004.	2.0E-004.
Body of mandible.	8.2E-004.	7.5E-004.
Angle of mandible.	9.3E-004.	8.2E-004.
Retromolar area	1.5E-003.	1.4E-003.
Anterior border of ramus.	1.4E-003.	7.1E-004.
Inferior border of mandible.	2.1E-003.	2.1E-003.

Table 4: Displacement of teeth in all three axis.

Teeth	Mesioangular Impaction			Horizontal Impaction.		
	X-axis.	Y-axis.	Z-axis.	X-axis.	Y-axis.	Z-axis.
Central Incisor	1.80E-07	-1.60E-07	-5.70E-08	1.80E-07	-1.40E-07	-4.80E-08
Lateral Incisor	2.00E-07	-2.60E-07	-8.90E-08	2.20E-07	-2.50E-07	-7.30E-08
Canine	2.80E-07	-4.00E-07	-4.00E-08	4.20E-07	-5.20E-07	-1.82E-07
1st Pre-Molar	5.30E-07	-7.00E-07	-2.10E-07	5.10E-07	-6.90E-07	-2.20E-07
2nd Pre Molar	9.50E-07	-9.00E-07	-1.20E-07	9.50E-07	-8.90E-07	-1.10E-07
1st Molar	1.60E-06	-1.30E-06	-2.70E-07	1.60E-06	-1.20E-06	6.80E-09
2nd Molar	2.10E-06	-1.60E-06	1.30E-07	2.00E-06	-1.60E-06	9.90E-08




Fig. 4: Tooth displacement in three planes of axis.




Fig. 5: Tooth displacement in Y and Z axis.

3.2. Tooth displacement

The present study also determines the amount of tooth displacement caused by mesio-angular and horizontally impacted 3rd molar on the adjacent teeth. The displacements in both the impactions were registered in sagittal plane (Y-axis), vertical plane (Z-axis), buccolingual

(X-axis) directions (Figure 4; Table 4).The displacement brought about by 10 grams of force for both mesio-angular and horizontal impaction caused mesial tilting of all the teeth i.e. from 2nd molar to central incisor; with the greatest tilting of 2nd molar. (Figure 5)

A line passing from the long axis of impacted third molar (mesio-angular and horizontal) and second molar is




Fig. 6: Mesioangular impaction- Angular changes before and after force application.




Fig. 7: Horizontal impaction- Angular changes before and after force application.




Fig. 8: Tooth displacement in X-axis.

constructed. The angle at the point of intersection of both the long axis was measured before and after the application of force. For mesio-angular impaction an increase of 0.027 degrees of angle, whereas in horizontal impaction a decrease of 0.025 degrees of angle is seen (Figures 6 and 7).

When seen along the vertical plane, there was extrusion seen with 2nd molar with mesio-angular impaction and extrusion in 2nd molar and 1st molar with horizontal impaction (Figure 5). And all the teeth exhibited buccal flaring in case of both impactions (Figure 8; Table 4).

4. Discussion

It is important to understand the stress distribution of erupting 3rd molar, as it helps in determining, the cause of dental crowding that has been a subject of controversy over the years. The use of FEA in this study has described, about the stress distribution in three-dimensional pattern. The study of tooth movements in all three planes is being introduced for the first time. Even the smallest changes are being detected using the finite element model, which was not possible in all the previous clinical studies.^{8–12}

In this study, it is observed that stress is distributed along the symphysis, body, and angle, inferior border of the mandible, anterior border of ramus and the retro molar area in both the model. Highest concentration of stress is present at the inferior border of the mandible (2.1E-3MPa) and lowest concentration of stress is at symphysis (2.0E-4MPa). However, when comparing the two models, higher areas of stresses are seen with mesio-angular impaction (Table 4) which might be due to a smaller area of contact as compared to horizontal impaction. This is valuable information as mesio-angular impaction, are the most common impaction.

The tooth displacements in both impactions are registered in sagittal, vertical and, buccolingual direction. According to Sakuda et al.,¹³ crowding can cause displacement at any interproximal contact, with the highest incidence of contact between cuspids and lateral incisors. Along the sagittal plane, greater tilting of 2nd molar is seen and gradually reducing to incisors, all the teeth exhibited mesial tilting, which is consistent with the finding of Richardson¹⁴ which states that late anterior crowding is associated with mesial movement of teeth rather than lower incisor retroclination. According to Sumitra et al.¹⁵ anterior component of occlusal force (ACF) is important in the discussion of late lower arch crowding. This is because it's possible role in causing mesial migration of teeth and subsequent dental mal-alignment. The axial inclination of permanent teeth is such that the forces of mastication produce a mesial resultant through the contact points of teeth. ACF is thought to result from axial inclination of posterior teeth which causes these teeth to tip forward during occlusal loading. This tendency of teeth to move forward varies greatly according to the angulations of teeth and by the steepness of occlusal plane.

In case of mesio-angular impaction there is a mesial inclination of the 3rd molar towards the long axis of 2nd molar at an angle of 0.027 degrees (Figure 6) whereas, in case of horizontal impaction the angle is 0.025 degrees (Figure 7), which is supported by study conducted by Artun et al.,¹⁶ which showed higher frequency of mesial angulation of 3rd molar in patients with impactions. But these studies are contrary to the study of Pirttiniemi et. al.,¹⁷ who concluded by his study that there is a slight distal drift of 2nd molar with no significant change in the lower incisor region. The present study describes about the stress

distribution in mesio-angular and horizontal impaction which was not mentioned in the previous studies.⁸⁻¹²

In vertical plane, there is extrusion with 2nd molar whereas all the other teeth exhibited intrusion in case of mesio-angular impaction, whereas in case of horizontal impaction extrusion is seen with 2nd molar and 1st molar and intrusion with all the other teeth. Along the buccolingual direction, in both the impaction all the teeth exhibited buccal flaring in which highest flaring is seen with 2nd molar (Table 4). This is similar to the result by Richardson who concluded that mesial inclination of lower canine is usually considered to be a sign of buccal segment moving forward. However, this finding is against the suggestion by Southard et al., who measured mesial force exerted by unerupted 3rd molar using a technique similar to measuring the anterior component of occlusal force. The mesial force increases tightness of all the proximal tooth contacts with no flaring and that surgical removal of 3rd molar will relieve the tightness by eliminating this force.

4.1. Limitations of the Study

FEM will give results with a greater degree of accuracy, but this approach has certain limitations, such as its inability to measure accurately the biological dynamics of the tooth and its supporting structure.¹⁸ The eruptive forces of the teeth are continuous process, however, the forces used in this study are not continuous, as it is not possible for the software to simulate the natural eruptive process. A common error in finite element models of bone in the literature is assigning one Young's modulus value to cortical bone and another to trabecular bone. Bone is a heterogeneous, anisotropic composite biomaterial which has variable Young's modulus based on its mineral content, thus the stress distribution in the present study may differ from the results obtained in the biologic subjects.

5. Conclusion

The results of finite element method simulations tested in the present study indicated that:

1. The magnitude of stress is more with mesio-angular than horizontal impaction.
2. High concentration of stress is present at the inferior border of the mandible whereas lowest concentration of stress is present at symphysis for both the models.
3. Along the sagittal axis, there is mesial tilting of all the teeth in both the impaction, which gradually decreased from 2nd molar to central incisor.
4. Along the vertical axis, in mesio-angular impaction there is extrusion of 2nd molar and intrusion with all the other teeth whereas, in horizontal impaction there is extrusion of 1st and 2nd molar and also intrusion of all the other teeth.

5. Along the bucco-lingual axis, buccal flaring of the teeth is seen.

From our finite element study data, it can be concluded that there are significant level of tooth displacement from 2nd molar towards the incisor after application of mandibular third molar eruptive force. These changes occurring along the dental arch must be taken into consideration while planning for prophylactic extraction of third molars.

6. Source of Funding

None.

7. Conflict of Interest

None.

References

1. Angelis MG, Blasi S, Pesce P, Lanteri V. Third molars and dental crowding: different opinions of orthodontists and oral surgeons among Italian practitioners. *Prog Orthod.* 2014;15(1):60. doi:10.1186/s40510-014-0060-y.
2. Bergstorm K, Jensen R. Responsibility of the third molar for secondary crowding. *Dent Abstr.* 1961;54:111–24.
3. Bishara S, Treder JE, Damon P, Olsen M. Changes in dental arches and dentition between 25 and 45 years of age. *Angle Orthod.* 1996;66(6):417–22. doi:10.1043/0003-3219(1996)066<0417:CITDAA>2.3.CO;2.
4. Vego L. A longitudinal study of mandibular arch perimeter. *Angle Orthod.* 1962;32(3):187–92. doi:10.1043/0003-3219(1962)032<0187:ALSOMA>2.0.CO;2.
5. Broadbent BH. Ontogenic development of occlusion. *Angle Orthod.* 1941;11(4):223–41. doi:10.1043/0003-3219(1941)011<0223:ODOO>2.0.CO;2.
6. Sidlauskas A, Trakiniene G. Effects of lower third molar on the lower dental arch crowding. *Stomatologija.* 2006;8(3):80–4.
7. Strang G, Fix G. An analysis of the finite element method. Englewood Cliffs (NJ: Prentice-Hall; 1973.
8. Kaplan RG. Mandibular third molars and post retention crowding. *Am J Orthod Dentofac Orthop.* 1974;66(4):411–30. doi:10.1016/0002-9416(74)90050-5.
9. Lindqvist B, Thilander B. Extraction of third molars in cases of anticipated crowding in the lower jaw. *Am J Orthod.* 1982;81(2):130–9. doi:10.1016/0002-9416(82)90037-9.
10. Richardson ME. The role of third molar in the cause of late lower arch crowding. A review. *Am J Orthod Dentofac Orthop.* 1989;95(1):79–83. doi:10.1016/0889-5406(89)90139-x.
11. Little RM. Stability and relapse of dental arch alignment. *Br J Orthod.* 1990;17(3):235–41. doi:10.1179/bjo.17.3.235.
12. Al-Balkhi KM. The effect of different lower third molar conditions on the re-crowding of lower anterior teeth in the absence of tight inter proximal contacts one year post orthodontic treatment: A Pilot study. *J Contemp Dent Pract.* 2004;15(3):66–73.
13. Sakuda M, Kuroda Y, Wada K, Matsumoto M. Changes in crowding of the teeth during adolescence and their relation to growth of the facial skeleton. *Trans Eur Orthod.* 1976;p. 93.
14. Richardson ME. Late lower arch crowding: facial growth or forward drift? *Eur J Orthod.* 1979;1(4):219–25. doi:10.1093/ejo/1.4.219-a.
15. Sumitra. Third Molars And Late Mandibular Incisor Crowding - A Review. *J Ind Orthod Soc.* 2005;38:100–11.
16. Artun J, Gard JD, Little RM. Long term stability of mandibular incisors following successful treatment of class II div I malocclusions. *Angle Orthod.* 1996;66(3):229–38. doi:10.1043/0003-3219(1996)066<0229:LTSOMI>2.3.CO;2.

17. Pirttiniemi PM, Oikarinen KS, Raustia AM. The effect of removal of third molars on the dental arches in the third decade of life. *Cranio*. 1996;12(1):23–7. doi:10.1080/08869634.1994.11677989.
18. Singh P, Wang C, Ajmera DH, Xiao SS, Song J, Ling Z, et al. Biomechanical Effects of Novel Osteotomy Approaches on Mandibular Expansion: 3D Finite Element Analysis. *J Oral Maxillofac Surg*. 2016;74(8):1658.e1–e15. doi:10.1016/j.joms.2016.04.006.

Meghna Bhandhari, Post Graduate

MD Baba Fareeduddin, Senior Lecturer

Cite this article: Kumar A, Bhandhari M, Baba Fareeduddin MD. The effects of erupting mandibular 3rd molar on the dental arch: A FEM study. *IP Indian J Orthod Dentofacial Res* 2022;8(3):173-179.

Author biography

Avinash Kumar, Professor